

# Lembo d'accesso mediante SPPT per trattamento endo-restaurativo di riassorbimento esterno

**Carmine Porciello**

*Private practitioner, Marigliano (Naples), Italy*

[SlidP – 2019 G. Vogel Award Session](#)

## **Clinical evaluation/ Diagnosis**

The patient was referred because of pain on a pink colouring of the cervical margin of tooth 1.1. We evaluated tooth vitality and took an x-ray, and we highlighted a root resorption. Probing depth was not pathological.

## **Treatment goals**

The treatment goals are the elimination of the pain, tooth survival, fisiological probing depth and good aesthetics.

## **Description of clinical/surgical procedures**

We used round burs to remove root demineralized tissues, so we verified that the resorption was in conjunction with pulp. We performed root canal therapy, then we executed an envelope flap through SPPT to expose root resorption and clean it. We positioned rubber dam and repaired the resorption with composite, then we removed periodontal attachment with the purpose of recreate a right biological width from the composite. We sutured the flap with polypropilene 6/0

## **Clinical outcomes**

After 10 days we removed sutures. The patient didn't use toothbrush and dental floss for 15 days. After 2 months we evaluated CAL loss and we found 1 mm REC and no pathological PD. At the 1 year control, all the treatment goals were respected although a 1 mm REC was noticed.